

Comment on “Quantum Phase Slips and Transport in Ultrathin Superconducting Wires”

In a recent Letter [1], Zaikin, Golubev, van Otterlo, and Zimanyi (ZGVZ) criticized the phenomenological time-dependent Ginzburg-Landau (TDGL) model which I used to study the quantum phase slip (QPS) for superconducting wires [2]. They claimed that they developed a “microscopic” model, made *qualitative* improvement on my overestimate of the tunnelling barrier due to electromagnetic field (EM), and claimed agreements with the experiments by Giordano [3].

In this comment, I want to point out that, i), ZGVZ’s result on EM barrier is expected in [2]; ii), their work is also phenomenological; iii), their renormalization scheme is fundamentally flawed; iv), they underestimated the barrier for ultrathin wires; v), their comparison with experiments is incorrect. Details are given below.

i), In [2] I emphasized results on relatively thick wires and concluded that the observations on Giordano’s wires with thickness $\sqrt{\sigma} = 410 - 505 \text{ \AA}$ must be due to weak links. Both kinetic inductance and Mooij-Schön mode have been included [2,4]. The ultrathin wire limit of the Eq.(8) in [2] gives an EM barrier,

$$\frac{S_{EM}}{\hbar} \cong \frac{\pi}{2} \sqrt{2 \ln(q_{upper}/q_{lower})} \frac{\sqrt{\sigma} \hbar c}{\lambda_L e^2}, \quad (1)$$

where the ratio of cutoffs $q_{upper}/q_{lower} \geq 10$. Eq. (1) is the main result in Ref. [1] except some underestimate (cf., below).

ii), QPS is a far-from-equilibrium process occurring on a time scale of the order of inverse gap of the superconductors. ZGVZ’s model contains nothing more than a saddle point plus Gaussian fluctuations and does not go beyond TDGL. Despite minor differences from my work, Ref. [1] does not present any new physics. I also want to point out that their electric screening for the superconducting charge is described by condensate fraction n_s which vanishes at T_c . This is incorrect since Debye screening is nondiscriminating against superconducting or normal charge.

iii), ZGVZ’s renormalization scheme is wrong since EM field *qualitatively* changes the Kosterlitz renormalization flow [4–6]. Therefore their conclusions on the “metal-superconductor” transition and wire resistance were not based on a solid foundation.

iv), ZGVZ underestimated the EM barrier for ultrathin wires. For Giordano’s wire of $\sigma = \pi(80 \text{ \AA})^2$, if we *assume* it is homogeneous with an effective London penetration depth $\lambda_L = 1000 \text{ \AA}$, the EM barrier *alone* reduces the tunneling probability to e^{-60} . Taking into account the substrate dielectric constant [4] $\epsilon = 10$ for *Ge*, then the EM suppression is e^{-330} . *In order to observe QPS in homogeneous wires, the wire radius must be smaller than $10 \text{ \AA}$.*

v), It needs to be emphasized that the results by Sharifi *et al.* [7] are NOT similar to that of [3], but qual-

itatively different. The wires in [7,8] are known to be homogeneous. In thinner wires ($[40 \text{ \AA}]^2$) than those in [3], no quantum phase-slip was observed. The wire resistance still obeys thermally activated behavior, albeit being larger than predicted by classical models [8]. As stated by ZGVZ, the theory is for a homogeneous wire. Hence they should compare with the experiments in [7] instead of [3]. A random Josephson junctions model [4] is suitable for granular wires [3,9].

TDGL was used to calculate the EM barrier for the quantum decay of supercurrent [2] and the EM contribution to vortex mass [10]. There is a deep physical reason [10] for the second order time derivative term in TDGL: superfluids are compressible due to number-phase conjugation. *Hence the qualitative physics in [2,10] is much more general than the formalism of the TDGL model.*

This work was supported by NSF (DMR 91-13631).

Ji-Min Duan

Department of Physics,
University of California-San Diego,
La Jolla, California 92093-0319

74.40.+k, 74.20.-z

-
- [1] A. D. Zaikin, D. S. Golubev, A. van Otterlo, and G. T. Zimanyi, Phys. Rev. Lett. **78**, 1552 (1997).
 - [2] J.-M. Duan, Phys. Rev. Lett. **74**, 5128 (1995).
 - [3] N. Giordano, Phys. Rev. Lett. **61**, 2137 (1988); Physica B **203**, 460 (1994).
 - [4] S. R. Renn and J.-M. Duan, Phys. Rev. Lett. **76**, 3400 (1996).
 - [5] M. P. A. Fisher and G. Grinstein, Phys. Rev. Lett. **60**, 208 (1988).
 - [6] S. C. Zhang, Phys. Rev. Lett. **59**, 2111 (1987).
 - [7] F. Sharifi, A. V. Herzog, and R. C. Dynes, Phys. Rev. Lett. **71**, 428 (1993).
 - [8] P. Xiong, A. V. Herzog, and R. C. Dynes, Phys. Rev. Lett. **78**, 927 (1997).
 - [9] A. V. Herzog, P. Xiong, F. Sharifi, and R. C. Dynes, Phys. Rev. Lett. **76**, 668 (1996).
 - [10] J.-M. Duan, Phys. Rev. B **48**, 333 (1993); *ibid*, **49**, 12381 (1994). J.-M. Duan, and A. J. Leggett, Phys. Rev. Lett. **68**, 1216 (1992).